INFOTRAC

- Help - Article
- Title List

- E-mail or Retrieval
- Links
- View mark list

Back to ...
- Citations
- Search
- Gale Group Databases

CIC University of Illinois Urbana Champ
Expanded Academic ASAP

Mark

Journal of Soil and Water Conservation, Wntr 2000 v55 i1 p79

No-tillage soybean performance in cover crops for weed management in the western Corn Belt. *M. M. Williams II; J. W. Doran; D. A. Mortensen.*

Full Text: COPYRIGHT 2000 Soil & Water Conservation Society

ABSTRACT: Frequency of soil moisture depletion and subsequent crop yield reduction resulting from water use by cover crops is unclear. Effects of cover crop residues and irrigation on soybean emergence, canopy volume, grain yield, and soil moisture in eastern Nebraska were determined Cover crop treatments included five common species and a bare soil control. Precipitation treatments included rainfed conditions and weekly irrigation events of 18 mm (0.7 in) in June and July. Above-normal May rainfall in 1995 and 1996 resulted in similar percent volumetric soil water contents (% VSWC) in the surface 15.2-cm (6.0-in) soil layer. Soybean emergence was unaffected by residue dry matter levels below 3,170 kg/ha (2,830 lb/ac). Early-season soybean growth was similar across all treatments; however, cover crop biomass greater than 2,170 kg/ha (1,940 lb/ac) reduced soybean canopy volume 33 to 44% during mid-season droughts. Yields were highest when stand densities were maintained and weeds were suppressed. This research develops a greater understanding of how cover crop residues influence soybean performance, ultimately reducing reliance on postemergent herbicide use for weed control in no-tillage systems.

Key words: Allelopathy, canopy volume, crop residues, fitness, integrated weed management, no-tillage, soil moisture, soil temperature, weather variability.

Cover crops are considered a tool in integrated weed management and control off-season soil and nutrient losses in row-crop production (Altieri and Liebman 1988; Swanton and Weise 1991). Residues on the soil surface with no-tillage management can reduce or delay crop and weed seedling recruitment, survival, and growth by altering the light environment, modifying soil temperatures, releasing allelochemicals, and acting as a physical barrier to emerging seedlings (Facelli and Pickett 1991). Allelochemicals have been isolated from commonly-studied cover crops, including barley (Hordeum vulgare L.), rye (Secale cereale L.), wheat (Triticum aestivum L.), and hairy vetch (Vicia villosa Roth.) (Shilling et al. 1985; White et al. 1989; Lovett and Hoult 1995). These residues inhibit growth of the foxtail species (Setaria spp.) and pigweed species (Amaranthus spp.) (Putnam et al. 1983; Mohler and Teasdale 1993). Several authors have demonstrated that weed suppression by cover crops complements effectiveness of reduced- rate herbicide programs in horticultural crops (Burgos and Talbert 1996; Wallace and Bellinder 1992).

Some studies have evaluated the suitability of cover crop systems for weed management in no-tillage soybean [Glycine max (L.) Merr.] production. In four years of a 5-yr study, soybean yielded significantly higher and weed biomass was reduced 74% to 91% in fall-planted rye compared to bare soil (Warnes et al. 1991). Improved soybean yields in two of three years corresponded with greater than 65% weed control from cover crop residues at 4 and 10 wk after planting (Liebl et al. 1992). Soybean yields in several winter cereal mulches were similar to yields in bare soil in four location-year experiments, except in one case where yields were improved by rye and triticale (X Triticosecale) residues that provided early-season weed suppression (Moore et al. 1994).

One concern about the use of cover crops in dryland agriculture is that these cover crops use soil moisture needed by the succeeding crop in dry years. Delaying the date of killing the cover crop

maximizes biomass production, assisting in weed suppression in the absence of tillage (Mohler and Teasdale 1993). However, allowing continued vegetative growth, particularly during a low-precipitation spring, may deplete stored soil moisture. In one of two years in Illinois, late-killed rye resulted in a 5-cm (2-in) loss of soil water (0-to-60 cm profile) in mid-June (Liebl et al. 1992). In Ontario, Wagner-Riddle et al. (1994) observed that rye killed 1 to 2 wk before planting reduced soil water content in one out of four sites/year combinations. Hairy vetch killed with a herbicide at corn planting in Nebraska had depleted soil moisture 3.1% compared to bare soil in the top 15 cm (5.9 in) in one out of four years (Power et al. 1991). Apparently, cover crops do not limit soil moisture for a succeeding crop in the Corn Belt every year. Less clear is the frequency of both soil moisture and crop yield reduction resulting from water use by cover crops.

Maintaining plant residues on the soil surface, as in the case of no-tillage crop production, can enhance soil moisture by decreasing runoff and evaporation (Adams et al. 1976). Liebl et al. (1992) found the upper 25 cm (9.8 in) of the soil profile had a higher %VSWC than bare soil for two rye kill dates when residues were left undisturbed on the soil surface. Likewise, Wagner-Riddle et al. (1994) saw an increase in soil water early in the soybean growing-season in one of four location-year experiments. In another study, rye mulch conserved soil moisture during drought periods of early corn and soybean growth, enabling greater plant water use deeper in the soil profile late in the season (Gallaher 1977).

Soil moisture is the most limiting resource in a rainfed environment; therefore, cover crops must be managed to minimize water stress on the subsequent crop. Although several studies have evaluated the impact of cover-crop mulches on soil water content in Illinois (Liebl et al. 1992) and Ontario (Wagner-Riddle et al. 1994), it is not known whether these results apply in the western Corn Belt where precipitation is more limited. Research indicates cover crops are valuable in Nebraska for soil stabilization (Walters 1987) and green manuring (Power et al. 1991). Cover crop growth characteristics and optimal planting dates have been identified for a sub-humid region (Power 1991; Power and Koerner 1994). However, few studies have characterized no-tillage soybean germination, growth, and yield in cover crop residues for weed management under contrasting soil moisture conditions.

The objectives of these field experiments were to identify how a cover crop mulch system for weed control influences soybean emergence, canopy growth, grain yield, and soil moisture under rainfed and irrigated conditions. The hypothesis being tested was that soybean has a unique set of environmental conditions (e.g., cover crop species and %VSWC) that optimize emergence, growth, and yield in cover crop residues.

Materials and methods

Site description. Field experiments were initiated in the fall of 1994 and 1995 at the University of Nebraska Agricultural Research and Development Center near Ithaca, Neb. Cover-crop treatments were planted on September 7 each year in fields where corn had been harvested for silage the preceding week (Table 1). The soil at both locations, 3.8 km (2.4 mi) apart, was a Sharpsburg silty clay loam (fine, montmorillonitic, mesic, typic Argiudoll). The soil at the first-year site contained 2.6% organic matter with a pH of 6.3, while the soil for the second-year site contained 2.8% organic matter and a pH of 5.8.

Experimental design and treatment structure. The experimental design was a randomized complete-block design with 3 replications in the fall of 1994, and 4 replications in 1995. Each block was randomized individually as a stripped split plot (split-block) with vertical strips of a "cover" treatment factor and (horizontal strips of a water treatment factor (horizontal strips orthogonal to vertical strips). "Cover" treatment levels in 1994 included a bare soil control plot, as well as barley (cv. Perkins), rye (cv. VNS), triticale (cv. Newcale), and wheat (cv. Arapaho), which were no-till drilled 2 cm (0.8 in) deep on 25-cm (9.8-in) rows in 31.0 m (100 ft) (15 ft x 4.6 in)) plots. Two additional cover crops were added in 1995: hairy vetch and a 1:1 ratio mixture of red clover (Trifolium repens L. cv. Mammoth) and yellow sweetclover [Melilotus officinalis (L.) Pall.]. Two "water" treatment levels included presence and absence of irrigation.

Cover crops were killed in the spring of each year (Table 1). In 1995, glyphosate [N-(phosphonomethyl) glycine] was applied over the entire experimental area at 2.2 kg ai/ha (2.0 lb/ac) with a tractor-mounted compressed-air sprayer at a carrier volume of 187 L/ha (20 gal/ac). In 1996, rye and wheat were killed with a hand-held [CO.sub.2] pressurized backpack sprayer at 1.1 kg ai/ha (1.0 lb/ac) of glyphosare and 187 L/ha (20 gal/ac) on May 13. Ten days later, the entire experimental site was sprayed with 1.4 kg ai/ha (1.3 lb/ac) of glyphosate. Since glyphosate had only a minor effect on hairy vetch, this cover crop species was clipped at the soil surface by hand within one week after herbicide application. Because of severe winterkill in the clover mixture, data were not collected from this treatment.

On June 7, 1995, and May 21, 1996, 250,900 seeds/ha (100,000 seeds/ac) of "Duhbar" soybean (group III indeterminate) were no-till planted 2 cm (0.8 in) deep in 76-cm (30-in) rows. A Buffalo-Till 4500-6AA (Fleischer Manufacturing, Columbus, Neb.) planter was used in 1995 and a Case 900 Cycloair (Case Corporation, Racine, Wisc.) planter the following year. A sprinkler irrigation system was assembled to irrigate randomly-assigned subplots measuring 7.6 x 4.6 m (25 x 15 ft). The irrigated plots received approximately 18 mm (0.7 in) of water applied weekly in June and July (Figures 1c and 2c).

Cover crop treatment provided the only means of weed suppression the first 5 wk after soybean planting. Grasses, predominantly foxtails (Setaria spp.), were controlled with 0.3 kg/ha (0.3 lb/ac) of sethoxydim [2-[1-(ethoxyimino)butyl] 5[2-(ethylthio)propyl]-3-hydroxy-2-cylohexen-1-one] plus recommended adjuvants applied with a tractor-mounted compressed-air sprayer on July 9, 1995, and July 2, 1996. Weeds other than redroot pigweed (Amaranthus retroflexus L.) and common waterhemp (Amaranthus rudis Sauer) were removed by hand.

Microsite conditions. Cover crop biomass was determined within 3 days of spraying by clipping the above-ground cover crop biomass in each of four 0.14 [m.sup.2] (1.5 [ft.sup.2]) quadrant frames per plot. Biomass samples were then ovendried and weighed.

Percent volumetric soil water content (%VSWC) estimates were based on soil sampling in both years. Methods are described in detail (Williams et al. 1998). Rain gauges in irrigated plots and weather data from the automated weather station at the research site provided precipitation amounts. The %VSWC was rarely measured immediately after irrigation; however, %VSWC at this time was estimated based on %VSWC measured prior to irrigation and the amount of water applied during irrigation (assumed 0% evaporatory loss of sprinkler irrigation water). Volumetric soil water content at 60% water-filled pore space (WFPS) was 31.0 at the 0-to-15.2 cm (6.0 in) soil depth, where WFPS = %VSWC [divided by] soil porosity, and soil porosity = (1 - soil bulk density + 2.65).

Soil maximum-minimum thermometers were used to monitor temperature fluctuations at a 1.3-cm (0.5-in) soil depth. Soil temperature was recorded and thermometers were reset approximately every third day during emergence. Mean soil temperature was calculated from maximum-minimum values.

Sampling and analytical procedures. Soybean seedling emergence in 1995 was monitored along 1 m (39 in) of crop row within each subplot. Emerged soybean seedlings were counted approximately every 5 days the first 4 wk after planting. Emergence was defined as the point at which cotyledons protruded above the soil. Sampling was expanded to include three 1-rn (39-in) lengths of crop row within each subplot in 1996.

Five soybean plants in 1995, and three in 1996, were randomly selected for nondestructive growth assessment approximately 2 wk after planting. Crop height was measured from the soil surface to the growing point, and maximum plant width was measured perpendicular to the crop row at 50% height. Based on measurements of row height and width, soybean canopy volume ([m.sup.3]/[m.sup.2]) was determined. Approximately 130 days after planting (DAP), grain yield was determined by hand-clipping three 1-m (39-in) lengths of crop row, and then threshing and

drying composite soybean seed to a constant weight.

Statistical analysis. Seedling emergence, canopy growth, and grain yield were analyzed with analysis of variance using the SAS MIXED model (SAS 1995). Main effects and interactions were examined and treatment comparisons were made with protected LSD tests at the 95% confidence level. Since there were significant year by treatment effect interactions, data were not pooled over years.

Results and discussion

Precipitation, cover-crop biomass, and soil water. A relatively wet spring and dry summer characterized both 1995 and 1996. The first year, precipitation in May was 144 mm (5.7 in), while June, July, and August were 33 mm (1.3 in), 21 mm (0.8 in), and 30 mm (1.2 in), respectively. The following year, 178 mm (7.0 in) of precipitation fell during May, while the next three months totaled 60 mm (2.4 in), 9 mm (0.4 in), and 25 mm (1.0 in), respectively. For comparison, the 30-yr average for these four months is 110 mm (4.3 in), 106 mm (4.2 in), 80 mm (3.1 in), and 92 mm (3.6 in), respectively. Due to the timing of rainfall events in the spring of 1995, glyphosate was applied to cover crops 3 wk later than planned. As a result, more than 3,100 kg/ha (2,800 lb/ac) of aboveground biomass was present in barley and more than 6,300 kg/ha (5,600 lb/ac) of rye, triticale, and wheat (Table 1). Dry conditions in the fall of 1995 and the spring of 1996 reduced survival of barley, triticale, and hairy vetch, resulting in 130 kg/ha (120 lb/ac), 750 kg/ha (670 lb/ac), and 650 kg/ha (580 lb/ac) of biomass, respectively. Of the species studied, rye and wheat survival and biomass production were consistently high in both years.

Throughout 1995, soils in the cover crop treatments and bare soil had similar %VSWC in the 0-to-15.2 cm (0-to-6.0 in) profile (Figure la, b). Likewise, in 1996, soils, rye, hairy vetch, and the control treatments had similar %VSWC with one exception; 15 DAP (June 5) when soils with rye had a higher %VSWC (Figure 2a, b). Although soil sampling in 1995 was limited to a depth of 15.2 cm (6.0 in), samples were taken to a depth of 30.5 cm (12.0 in) the following year and no apparent differences in the %VSWC at the 15.2-to-30.5 cm (6.0-to-12.0 in) depth were detected. Apparently, soil moisture used by cover crops during growth was replenished by spring precipitation in both years.

Irrigation in June and July of both years increased %VSWC relative to the rainfed treatment (Figure la, b; Figure 2a, b). Lack of precipitation events in June and July frequently reduced the 0-to-15.2 cm (0-ro-6.0 cm) soil profile below PWP.

Soybean emergence. The presence of all residues suppressed soybean emergence early and reduced final soybean plant densities 34% (barley) to 60% (rye) (Figure 3a). Poor seedling emergence in cover crop residues in 1995 was likely the result of insufficient soil/seed contact. Because most residues exceeded 6,310 kg/ha (5,630 lb/ac), an unusually large amount of residue was left on the soil surface. Slightly higher soil moisture content in residue treatments observed at soybean planting further complicated planting operations and the front coulter of the planter had difficulty cutting through cover crop residues. As a result, residue was pressed into a moist furrow, preventing the furrow from closing. High residue levels have interfered with soybean planting in other studies, particularly where planting equipment failed to penetrate surface residues (Eckert 1988; Hovermale et al. 1979; Liebl et al. 1992; WagnerRiddle et al. 1994).

Conditions were drier at planting in 1996, and lower amounts of cover crop residue [130 to 2,890 kg/ha (120 to 2,580 lb/ac)] resulted in more effective seed placement. With the exception of wheat residue at 11 DAP, cover crop residues did not delay soybean emergence, and final soybean plant densities were similar among cover crops (Figure 3b).

Soybean growth. Measurement of plant volume as an indicator of size provides a non-destructive index of plant weight, and in the case of many annual species, seed production (Bussler et al. 1995). Canopy volume, as it was measured in this study, characterized the space occupied by the above ground portion of the soybean row. In making comparisons across treatments, plant

biomass per unit of canopy volume is assumed constant.

The presence of cover crop residues in 1995 reduced mid-season soybean canopy volume. Early in the season, canopy volumes were similar in all cover-crop residues, yet by 50 DAP, canopy volumes in all residue treatments were less than the bare soil control (36% less with wheat, and 29% less with triticale) (Figure 4a). The 1995 canopy volume measurements probably reflect the soybean plant density differences in which the presence of residues resulted in 34 to 60% fewer seedlings. Reduced plant densities may have altered morphology of individual plants, with less vertical growth and greater between-plant growth in residue treatments. This would explain similar per plant soybean grain yield (Table 2) in all cover crop residues, ranging from 12.3 g/plant (0.43 oz/plant) in barley, to 15.0 g/plant (0.53 oz/plant) in wheat. Therefore, while a lower seedling population in cover-crop residues reduced soybean canopy growth, individuals in all treatments had the same fitness at the end of the season.

Soil moisture limited soybean canopy volume in 1995 (Figure 4b). June and July precipitation was 52 mm (2.0 in), considerably less than the 30-yr average of 189 mm (7.4 in). Although soil moisture was slightly lower in the control plots at 35 DAP (July 12), the fact that %VSWC was below the PWP indicates plants obtained adequate soil moisture from below the 15.2-cm (6.0-in) soil depth to sustain soybean growth during moisture stress periods (Figure la, b). On several occasions, more soil moisture was available in irrigated plots. Consequently, canopy volume was greater in irrigated plots beyond 50 DAP (Figure 4b).

In 1996, the presence of rye and wheat residues delayed initial soybean canopy development. The presence of rye and wheat residues reduced mid-season canopy volume 50 and 54%, respectively (Figure 5). Stone and Taylor (1983) found taproot and lateral root extension increases with soil temperature. Rye residue resulted in a lower rate of soil warming. For instance, rye treatments took approximately 2 wk longer for mean soil temperature at the 1.3-cm (0.5-in) depth to reach 26 [degrees]C (79[degrees] F) (Figure 6b). It is plausible that the delay in soil warming through 36 DAP delayed soybean canopy growth through 63 DAP.

Reduced early-season soybean canopy volume in 1996, as well as emergence, may also have been due in part to allelopathic compounds originating from decaying cover crop residue. Primary compounds and metabolires inhibiting germination and growth of a number of plant species have been isolated from barley, rye, wheat, and hairy vetch (Shilling et al. 1985; White et al. 1989; Lovett and Hoult 1995). In addition, frequent irrigation and natural precipitation events may have enhanced conditions favoring anaerobic microbial activity under high residue levels (Williams et al. 1998). As a result, phytotoxic compound production may have occurred several weeks after cover crop death. For instance, acetic acid is released from decomposing wheat straw (Lynch 1977).

The presence of cover crop residues resulted in greater canopy size later in the 1996 season due in part to weed suppression. At 3 wk after planting, Amaranthus spp. canopy volume was reduced 38, 71, 48, 58, and 67% by presence of barley, rye, triticale, wheat, and hairy vetch residues, respectively (Williams et al. 1998). With respect to the bare soil treatment, soybean canopy volume increased 16, 27, and 28% in barley, wheat, and hairy vetch residues at 79 DAP, respectively (Figure 5). In addition, individual soybean grain yield was 9.2 g/plant (0.32 oz/plant), 9.1 g/plant (0.32 oz/plant), and 7.4 g/plant (0.26 oz/plant) for rye, wheat, and hairy vetch residues (Table 2), on average 54% higher than the control and indicating a larger canopy biomass by harvest in these residues. Apparently, early-season suppression of weeds by the presence of cover-crop residues reduced weed competitiveness and increased soybean fitness.

Soybean grain yield. An interaction between cover crop and water on soybean grain yield was observed in 1995 (Table 3). Under irrigation, with the exception of the presence of rye, which reduced yields 11%, soybean yields in the bare soil and cover crops were not significantly different. However, in the absence of supplemental irrigation water, the presence of residues resulted in significantly lower soybean yields. Since per plant grain yields were identical across all cover treatments (Table 2), differences in total yields were a function of soybean density. As

discussed earlier, interference with planting operations from crop residue has lead to decreased soybean yields in other investigations (Eckert 1988; Liebl et al. 1992).

In 1996 the presence of rye, wheat, and vetch residues resulted in higher soybean grain yields compared to the baresoil treatment (Table 2). Despite the slight delay in soybean emergence in the presence of wheat residue and mid-season stunting by the presence of rye and wheat residue, soybean in those environments yielded 990 kg/ha (880 lb/ac) and 760 kg/ha (680 lb/ac) more than the bare soil control. As mentioned earlier, suppression of weeds by the presence of cover crop residues early in the season may have reduced their competitive effect on the crop. When cover crop residues minimized weed competition, other studies have found enhanced crop yields (Warnes et al. 1991; Liebl er al. 1992; Moore et al. 1994).

Conclusions

Winter cover crops play a role in soil and water conservation and integrated weed management in soybean throughout the Corn Belt. They also can be used in semiarid regions provided the dynamic resource needs of the crop are met. Our findings in eastern Nebraska indicate no-tillage soybean can yield well in residues when cover crop residues do not reduce soybean stand densities. Soybean emergence declined as cover crop biomass rates exceeded approximately 3,100 kg/ha (2,800 lb/ac). Williams et al. (1998) found Amaranthus spp. and Setaria spp. growth are reduced 37% to 97% at 3 to 5 wk after planting into cover crop residues with biomass levels as low as 130 kg/ha (120 lb/ac). Furthermore, early-season soybean growth is frequently placed at a competitive advantage over that of the weeds.

In both years of this research, above-normal precipitation in May was followed by below-normal precipitation in June, July, and August. Transpirational demands from cover crops coincided with high precipitation events, explaining why %VSWC in cover-crop treatments were never lower than the control.

Additional research needs to look more closely at year-to-year variability in weather patterns and how they influence cover-crop water usage and soybean performance under dryland conditions. Ultimately, coupling such modeling efforts with recent findings on weed suppression will provide the necessary framework for establishing guidelines that minimize yield-reducing risks of using cover crops, as well as maintain the crop at a competitive advantage over the weeds. Exploiting crop and weed selectivity to cover crops offers an approach to conserve both soil and water resources in no-tillage systems that historically rely heavily upon herbicides for weed control.

Martin M. Williams II is graduate research assistant, and David A. Mortensen is professor, Department of Agronomy, University of Nebraska-Lincoln, Lincoln, Neb.; and John W Doran is soil scientist with the US. Department of Agriculture's Agricultural Research Service (USDA-ARS), Lincoln, Neb. They thank R. Drijber, J. Lindquist, and A. Martin for discussion, and E. Anderson, G. Lesoing, and D. Radach for technical assistance, This research was a joint project of the University of Nebraska Agricultural Research Division and the USDA-ARS, Journal series no. 12043.

REFERENCES CITED

Adams, J.E., G.F. Arkin, and J.T. Ritchie. 1976. Influence of row spacing and straw mulch on first stage drying. Journal of Soil Science 40:436-442.

Altieri, M.A., and M. Liebman. 1988. Weed management: Ecological guidelines. In: M.A. Altieri and M. Liebman (Eds.) Weed Management in Agroecosystems: Ecological Guidelines. CRC Press, Inc.: Boca Raton, FL.

Burgos, N.R., and R.E. Talbert. 1996. Weed control and sweet corn (Zea mays var. Rugosa) response in no-till system with cover crops. Weed Science 44:355-361.

Bussler, B.H., B.D. Maxwell, and K.J. Puettmann. 1995. Using plant volume to quantify interference in corn (Zea mays) neighborhoods. Weed Science 43:586-594.

Eckert, D.J. 1988. Rye cover crops for no-tillage corn and soybean production. Journal of Production Agriculture 1:207-210.

Facelli, J.M., and S.T.A. Pickett, 1991. Plant litter: Its dynamics and effects on plant community structure. Botanical Review 57:1-32.

Gallaher, R.N. 1977. Soil moisture conservation and yield of crops no-till planted in rye. Journal of Soil Science 41:145-147.

Hovermale, C.H., H.M. Camper, and M.W. Alexander. 1979. Effects of small grain stubble height and mulch on no-tillage soybean production. Agronomy Journal 71:644-647.

Liebl, R., F.W. Simmons, W.M. Wax, and E.W. Stoller. 1992. Effect of rye (Secale cereale) mulch on weed control and soil moisture in soybean (Glycine max). Weed Technology 838-846.

Lovett. J.V., and A.H.C. Hoult. 1995. Allelopathy and self-defense in barley. In: Inderjit, K.M.M. Dakshini, and F.A. Einhellig (Eds.). Allelopathy: Organisms, Processes, and Applications. American Chemical Society: Washington, DC.

Lynch, J.M. 1977. Phytotoxicity of acetic acid produced in the anaerobic decomposition of wheat straw. Applied Bacteriology 42:81-87.

Mohlder, C.L., and J.R. Teasdale. 1993. Response of weed emergence to rate of Vicia villosa Roth and Secale cereale L. residue. Weed Research 33:487-499.

Moore, M.J., T.J. Gillespie, and C.J. Swanton. 1994. Effect of cover crop mulches on weed emergence, weed biomass, and soybean (Glycine max) development. Weed Technology 8:512-518.

Power, J.F. 1991. Growth characteristics of legume cover crops in a semiarid environment. Journal of Soil Science 55:1659-1663.

Power, J.F., J.W. Doran, and P.T. Koerner. 1991. Hairy vetch as a winter cover crop for dryland corn production. Journal of Production Agriculture 4:62-67.

Power, J.F., and P.T. Koerner. 1994. Cover crop production for several planting dates in eastern Nebraska. Agronomy Journal 86:1092-1097.

Putnam, A.R., J. DeFrank, and J.P. Barnes. 1983. Exploitation of alleloparhy for weed control in annual and perennial cropping systems. Chemical Ecology 9(8):1001-1010.

SAS Institute. 1995. SAS User's Guide, SAS Institute: Cary, NC.

Shilling, D.G., R.A. Liebl, and A.D. Worsham. 1985. Rye (Secale cereale L.) and wheat (Triticum aestivum L.) mulch: The suppression of certain broadleaved weeds and the isolation and identification of phytotoxins. In: A.C. Thompson (Ed.) The Chemistry of Allelopathy. American Chemical Society: Washington, DC.

Stone, J.A., and H.M. Taylor. 1983. Temperature and the development of the taproot and lateral roots of four indeterminate soybean cultivars. Agronomy Journal 75:613-618.

Swanron, C.J., and S.F. Weise. 1991. Integrated weed management: the rationale and approach. Weed Technology 5:657-663.

Wagner-Riddle, C., T.J. Gillespie, and C.J. Swanron. 1994. Rye cover crop management impact on soil water content, soil temperature and soybean growth. Canadian Journal of Plant Science 74:485-495.

Wallace, R.W., and R.R. Bellinder. 1992. Alternative tillage and herbicide options for successful weed control in vegetables. HortScience 27:745-749.

Walters, D.T. 1987. Early studies on the use of legumes for conservation tillage in Nebraska. In: J.F. Power (Ed.) Cover Crops for Clean Water. Soil and Water Conservation Society: Ankeny, IA.

Warnes, D.D., J.H. Ford, C.V. Eberlein, and W.E. Lueschen. 1991. Effects of winter rye cover crop system and available soil water on weed control and yield in soybeans. In: W.L. Hargrove (Ed.) Cover Crops for Clean Water. Soil and Water Conservation Society: Ankeny, IA.

White, R.H., A.D. Worsham, and U. Blum. 1989. Allelopathic potential of legume debris and aqueous extracts. Weed Science 37:674-679.

Williams, M.M. II, D.A. Mortensen, and J.W. Doran. 1998. Assessment of weed and crop fitness in cover crop residues for integrated weed management. Weed Science 46:595-603.

Cover crop management.

Cover crop	Year 1995 Seeding rate kg/ha	Spray date	Biomass kg/ha	1996 Seeding rate kg/ha	Spray date	Biomass kg/ha
Barley	78.4	6 June 1995	3170	78.4	23 May 1996	130
Rye	78.4	6 June 1995	6310	78.4	13 May 1996	2890
Triticale	156.4	6 June 1995	7160	78.4	13 May 1996	750
Wheat	78.4	6 June 1995	6710	78.4	23 May 1996	2170
Vetch	--	--	--	35.8	23 May 1996	650
LSD (0.05)			780			330

Effects of cover and water treatments on no-tillage soybean grain yield.

Treatment	Soybean yield Individual 1995	1996	Total 1995	1996
	g/plant		kg/ha	
Cover				
Control	13.9	5.6	2060	1200
Barley	12.3	6.1	1630	1480
Rye	12.8	9.2	1540	2190
Triticale	13.2	6.0	1800	1440
Wheat	15.0	9.1	1920	1960
Vetch	--	7.4	--	1700
LSD (0.05)	(NS)	(2.7)	(230)	(630)
Water				
Irrigated	16.0	7.1	2150	1610
Rainfed	10.9	7.4	1430	1710
	(*)	(NS)	(*)	(NS)
Interactions				
Cover/water	NS	NS	*	NS

(*.)Significant at the 0.05 probability level.

Effects of cover and water treatments on no-tillage soybean grain yield in 1995.

Yield

Cover	Irrigated	Rainfed
	kg/ha	
Control	2150ab	1980a
Barley	2160ab	1100c
Rye	1910b	1170c
Triticale	2200ab	1410bc
Wheat	2330a	1510b
LSD (0.05)	350	

Within columns, means followed by the same letter are not significantly different at LSD (0.05).

Article A60013070

View other articles linked to these subjects:

Allelopathy - Evaluation
View 2 Periodical references
See also 7 other subdivisions
Cover Crops - Evaluation
See also 12 other subdivisions
No-Tillage - Management
View 8 Periodical references
See also 17 other subdivisions
Soil Conservation - Management
View 48 Periodical references
See also 61 other subdivisions
Soybean Industry - Management
View 37 Periodical references
See also 98 other subdivisions
Weeds - Control
View 270 Periodical references
See also 36 other subdivisions

Journal of Soil and Water Conservation, Jan 1, 2000
View other articles in this issue

Print, e-mail, and other retrieval options

Browser Print — *Full Text* —
Reformat for printing (approximately 11 pages) from your browser. To return to InfoTrac, use the *back* function of your browser.

Acrobat Reader — *Full Text* —
Retrieve for viewing and printing from Acrobat™ Reader. Please allow a few minutes for the retrieval operation to complete (approximately 7 pages)

E-Mail Delivery — *Full Text* —

We will send a plain text version to the e-mail address you enter (e.g. *bettyg@library.com*).

E-Mail Address:

Subject (defaults to title):

Submit E-mail Request

PostScript — *Full Text* —
Download article in PostScript format.

InfoTrac Articles — *Full Text* —

We will print this item at a nearby InfoTrac Articles Station **(Locations)**. Use the order name you enter here to identify your print job.

Order Name:

Submit Print Order

—— **Article 1 of 1** ——

THOMSON GALE

Copyright and Terms of Use